arXiv:1603.00830v2 [math.DS] 9 Mar 2016

LOEWNER EVOLUTION OF HEDGEHOGS AND 2-CONFORMAL MEASURES OF CIRCLE MAPS

KINGSHOOK BISWAS

Abstract. Let f be a germ of holomorphic diffeomorphism with an irrationally indifferent fixed point at the origin in $\mathbb{C}$ (i.e. $f(0) = 0, f'(0) = e^{2\pi i\alpha}, \alpha \in \mathbb{R} - \mathbb{Q}$). Perez-Marco (Fixed points and circle maps, Acta Math. **179** (2) 1997, 243-294) showed the existence of a unique continuous monotone one-parameter family of nontrivial invariant full continua containing the fixed point called Siegel compacta, and gave a correspondence between germs and families (g_t) of circle maps obtained by conformally mapping the complement of these compacts to the complement of the unit disk. The family of circle maps (g_t) is the orbit of a locally-defined semigroup (Φ_t) on the space of analytic circle maps which we show has a well-defined infinitesimal generator X. The explicit form of X is obtained by using the Loewner equation associated to the family of hulls (K_t). We show that the Loewner measures (μ_t) driving the equation are 2-conformal measures on the circle for the circle maps $(z \mapsto \overline{g_t(\overline{z})})$.

AMS Subject Classification: 37F50

Contents

1. Introduction 1
2. Boundary values of the Herglotz transform 5
3. The Loewner equation 6
4. Conformal measures of analytic circle diffeomorphisms 7
5. The semigroup $(\Phi_t)_{t\geq 0}$ on $\mathrm{Diff}^{\omega}_{\alpha}(S^1)$ 7
6. Infinitesimal generator of the semigroup 8
7. Proofs of main results 10
8. Linearizable maps and conformal radius of linearization domains 13
References 16

1. Introduction

A germ $f(z) = e^{2\pi i\alpha}z + O(z^2), \alpha \in \mathbb{R} - \mathbb{Q}/\mathbb{Z}$ of holomorphic diffeomorphism fixing the origin in $\mathbb{C}$ is said to be linearizable if it is analytically conjugate to the rigid rotation $R_\alpha(z) = e^{2\pi i\alpha}z$. The number α is called the rotation number of f, and the maximal domain of linearization is called the Siegel disk of f. The linearizability of f is dependent on the arithmetic of α, and the optimal arithmetic condition for linearizability in this setting is given by the well-known Brjuno condition (see [Sie42], [Brj71], [Yoc95]).

Perez-Marco proved the existence of a unique, strictly increasing Hausdorff continuous family $(K_t)_{t>0}$ of non-trivial invariant full continua containing the fixed point called *Siegel compacts* ([PM97]), where $K_t \to \{0\}$ as $t \to 0$, and K_t can be described as the connected component containing the origin of the set of non-escaping points in the closed disc $\overline{\mathbb{D}}_t$ of radius t around the origin. When f is non-linearizable these are called *hedgehogs*. The topology and dynamics of hedgehogs has been studied by Perez-Marco ([PM94], [PM96]), who also developed techniques using "tube-log Riemann surfaces" ([BPM15a], [BPM15b], [BPM13]) for the construction of interesting examples [PM93], [PM95], [PM00] of indifferent germs and hedgehogs, which were also used by the author ([Bis05], [Bis08], [Bis15]) and Cheritat ([Che11]) to construct further examples.

Notation: Throughout $\mathbb{D}_\infty$ will denote the complement in $\hat{\mathbb{C}}$ of the closed unit disk, $\mathbb{D}_\infty = \hat{\mathbb{C}} - \overline{\mathbb{D}}$, $r : S^1 \to S^1$ will denote complex conjugation, $r(\xi) = \overline{\xi}$, and $\frac{d^+}{dt}$ will denote right-hand derivative.

The construction of Perez-Marco from [PM97] associates to a pair (f, K) an analytic circle diffeomorphism g, where f is a germ with an irrationally indifferent fixed point at the origin and K is a Siegel compact of f, by considering a conformal map ψ from the complement of K to the complement of the closed unit disk, $\psi : \hat{\mathbb{C}} - K \to \mathbb{D}_\infty$, such that $\psi(\infty) = \infty$. Conjugating f by ψ gives a holomorphic diffeomorphism $g = \psi \circ f \circ \psi^{-1}$ in an annulus in $\mathbb{D}_\infty$ having S^1 as a boundary component, and g is shown to extend across S^1 to an analytic circle diffeomorphism defined in a neighbourhood of S^1 such that the rotation numbers of g and f are equal, $\rho(g) = \rho(f) = \alpha$.

Invariant compacts for g containing S^1 then correspond to invariant compacts for f containing K, and the theorem on existence and uniqueness of Siegel compacts then gives the existence and uniqueness of a continuous strictly increasing 1-parameter family of *Herman compacts* $(A_t)_{0 \leq t < \epsilon}$ for the circle map g with $A_0 = S^1$, where a Herman compact for g is a connected totally invariant compact A containing S^1 such that $\hat{\mathbb{C}} - A$ has two components (such a compact is necessarily invariant under reflection in the unit circle as well). The construction which produces g from the pair (f, K) can similarly be applied to each pair (g, A_t) to give a one-parameter family of analytic circle diffeomorphisms $(g_t)_{t \geq 0}$ with $g_0 = g$, by conjugating g by a conformal map $\psi_t : \hat{\mathbb{C}} - (\overline{\mathbb{D}} \cup A_t) \to \mathbb{D}_\infty$, normalized so that $\psi_t(\infty) = \infty, \psi_t'(\infty) > 0$. As before the map $g_t = \psi_t \circ g \circ \psi_t^{-1}$ defined in a one-sided neighbourhood of S^1 extends across to give an analytic circle diffeomorphism g_t with rotation number equal to that of g.

The family A_t being Hausdorff continuous and strictly increasing, it is possible to reparametrize the family so that the compacts $\overline{\mathbb{D}} \cup A_t$ have logarithmic capacity t, meaning that the expansion of ψ_t near $z = \infty$ is of the form

$$\psi_t(z) = e^{-t} z + a_0(t) + \frac{a_1(t)}{z} + \dots$$

Let $\mathrm{Diff}^\omega(S^1)$ and $\mathrm{Diff}^\omega_\alpha(S^1)$ denote the space of analytic circle diffeomorphisms and the subspace of those diffeomorphisms with fixed rotation number $\alpha \in (\mathbb{R} - \mathbb{Q})/\mathbb{Z}$ respectively. We obtain for $t \geq 0$ a locally defined family of maps $\Phi_t : \mathcal{D}_t \subset \mathrm{Diff}^\omega_\alpha(S^1) \to \mathrm{Diff}^\omega_\alpha(S^1)$ mapping a circle map g to the circle map g_t obtained from the above construction. Here the domain $\mathcal{D}_t$ of Φ_t consists of those circle maps in

$\mathrm{Diff}^{\omega}_{\alpha}(S^1)$ which have a Herman compact A such that $\overline{\mathbb{D}} \cup A$ has logarithmic capacity t. It is not hard to show that in fact the maps Φ_t form a continuous semigroup, namely $\Phi_0 = id$,

$$\Phi_s \circ \Phi_t = \Phi_{s+t}$$

whenever the composition is defined, and the orbits $t \mapsto \Phi_t(g)$ give continuous curves in $\mathrm{Diff}^{\omega}_{\alpha}(S^1)$.

We show in fact that the semigroup $(\Phi_t)_{t \geq 0}$ has an infinitesimal generator X, meaning the curves $t \mapsto \Phi_t(g)$ are right-hand differentiable in t and

$$\frac{d^+}{dt}\Phi_t(g) = X(\Phi_t(g))$$

Since the space $\mathrm{Diff}^{\omega}_{\alpha}(S^1)$ does not carry any obvious differentiable structure, the sense in which these assertions hold is made precise in the statements of the theorems below.

The form of the infinitesimal generator X is obtained by studying the *Loewner equation* associated to the family of hulls $\overline{\mathbb{D}} \cup A_t$ (we recall in section 2 the basic facts about the Loewner equation which we will be needing). The maps $\phi_t := \psi_t^{-1} : \mathbb{D}_\infty \to \hat{\mathbb{C}} - (\overline{\mathbb{D}} \cup A_t)$ form a *Loewner chain*, and it is known that $t \mapsto \phi_t(z,t)$ is absolutely continuous for each fixed $z \in \mathbb{D}_\infty$. Moreover, for almost every t, the right-hand derivatives

$$\chi_t(z) = \frac{d^+}{ds}_{|s=t}(\phi_t^{-1} \circ \phi_s)(z)$$

exist and the functions $H_t(z) = \chi_t(z)/z$ are given by the Herglotz transforms on $\mathbb{D}_\infty$ of a family of probability measures on the unit circle (μ_t), called the *driving* or *Loewner measures* of the Loewner equation. Here by the Herglotz transform on $\mathbb{D}_\infty$ of a probability measure μ on S^1 we mean the holomorphic function $H = \mathcal{H}\mu$ in $\mathbb{D}_\infty$ with positive real part and satisfying $H(\infty) = 1$, defined by

$$(\mathcal{H}\mu)(z) = \int_{S^1} \frac{\xi + 1/z}{\xi - 1/z} d\mu(\xi)$$

(the classical Herglotz Theorem asserts that any holomorphic function H on $\mathbb{D}_\infty$ satisfying $\mathrm{Re}\, H > 0, H(\infty) = 1$ is of this form for a unique probability measure μ on S^1). Douady and Yoccoz have shown in [DY99] the existence and uniqueness of an s-conformal measure $\mu = \mu_{s,g}$ for any C^2 circle diffeomorphism g with irrational rotation number and any $s \in \mathbb{R}$. By an s-conformal measure for a circle diffeomorphism g we mean a probability measure μ on S^1 such that

$$\mu(g(E)) = \int_E |g'(x)|^s d\mu(x)$$

for all measurable sets $E \subset S^1$.

We show that for any g in $\mathrm{Diff}^{\omega}_{\alpha}(S^1)$, the associated Loewner chain (ϕ_t) is differentiable for **all** t uniformly for z in any compact subset of $\mathbb{D}_\infty$, and the associated Loewner measures are the pull-backs by complex conjugation of the 2-conformal measures of the circle diffeomorphisms $g_t = \Phi_t(g)$:

Theorem 1.1. *(Loewner measures are 2-conformal measures). For any g in $Diff^{\omega}_{\alpha}(S^1)$ and any $t \geq 0$, the right-hand derivative of the Loewner chain (ϕ_t) exists (uniformly for z in any compact subset of $\mathbb{D}_{\infty}$) and is given by*

$$\chi_t(z) = \frac{d^+}{ds}_{|s=t}(\phi_t^{-1} \circ \phi_s)(z) = z \cdot (\mathcal{H}r^* \mu_{2,g_t})(z)$$

where μ_{2,g_t} is the unique 2-conformal measure of g_t.

The existence and form of the infinitesimal generator X of the semigroup (Φ_t) is stated as follows:

Theorem 1.2. *(Infinitesimal generator of the semigroup). For any $g \in Diff^{\omega}_{\alpha}(S^1)$, there exists a function $X(g)$ holomorphic in a neighbourhood V of S^1 such that for $t > 0$ small the circle maps $\Phi_t(g)$ are defined in V and*

$$\lim_{t \to 0^+} \frac{\Phi_t(g) - g}{t} = X(g)$$

uniformly on compacts in V. The holomorphic function $X(g)$ is given (in $V \cap \mathbb{D}_{\infty}$) by

$$X(g) = g' \cdot \chi - \chi \circ g$$

where $\chi(z) = z \cdot (\mathcal{H}r^* \mu_{2,g})(z)$.

We can think of the curves $(t \mapsto \Phi_t(g))$ as integral curves of a vector field X on the space $\mathrm{Diff}^{\omega}_{\alpha}(S^1)$. The next theorem asserts the uniqueness in forward time for these integral curves:

Theorem 1.3. *(Uniqueness in forward time of integral curves). If $(g_t)_{0 \leq t < \epsilon} \subset Diff^{\omega}_{\alpha}(S^1)$ is a family of circle diffeomorphisms holomorphic in a neighbourhood V of S^1 such that the right-hand derivatives exist and satisfy*

$$\frac{d^+}{ds}_{|s=t} g_s(z) = X(g_t)(z)$$

uniformly on compacts in V, then $g_t = \Phi_t(g_0)$ for $0 \leq t < \epsilon$.

Definition 1.4 (Germ of integral curve)**.** *An integral curve (in the sense of the previous theorem) $(g_t)_{-\infty < t < t_0} \subset Diff^{\omega}_{\alpha}(S^1)$ defined for all times t less than some t_0 is said to be a backward integral curve of X. We say that two backward integral curves $(g^1_t)_{-\infty<t<t_1}, (g^2_t)_{-\infty<t<t_2}$ of X define the same germ of integral curve of X if $g^1_t = g^2_t$ for all $t < t_0$ for some $t_0 < t_1, t_2$.*

It follows from the above that if $(g^1_t)_{-\infty<t<t_0}, (g^2_t)_{-\infty<t<t_0}$ are two backward integral curves defining the same germ of integral curve then in fact $g^1_t = g^2_t$ for **all** $t < t_0$. Any germ f with an irrationally indifferent fixed point gives a family of Siegel compacts $(K_t)_{-\infty<t<t_0}$ (parametrizing the compacts K_t by their logarithmic capacities), and hence (by applying the germs to circle maps construction to the pairs (f, K_t)) gives a family of circle maps $(g_t = g^f_t)_{-\infty<t<t_0}$ which it is easy to see is a backward integral curve. We show that conversely any backward integral curve arises in this way from a germ f. Denoting the space of germs with rotation number α by $\mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$ we have:

Theorem 1.5. *(Germs of diffeomorphisms and germs of integral curves). For any backward integral curve $(g_t)_{-\infty<t<t_0} \subset Diff^{\omega}_{\alpha}(S^1)$ of X, we have $g_t \to R_\alpha$ uniformly on S^1 as $t \to -\infty$.*
The map $f \mapsto [(g_t^f)]$ gives a one-to-one correspondence between $Diff_\alpha(\mathbb{C}, 0)$ and germs of integral curves of X.

Finally in section 8 we describe some further results that can be obtained in the case of analytically linearizable circle maps and germs.

Acknowledgements The author thanks Ricardo Perez-Marco for helpful discussions.

2. Boundary values of the Herglotz transform

The Herglotz transform of a probability measure μ on S^1 is the holomorphic function $\mathcal{H}\mu$ in $\mathbb{D}$ defined by

$$(\mathcal{H}\mu)(w) = \int_{S^1} \frac{\xi + w}{\xi - w} d\mu(\xi)$$

The real and imaginary parts of the Herglotz transform $\mathcal{H}\mu = \mathcal{P}\mu + i\mathcal{Q}\mu$ are given by the Poisson and conjugate Poisson transforms,

$$(\mathcal{P}\mu)(w) = \int_{S^1} \mathrm{Re}\, \frac{\xi + w}{\xi - w} d\mu(w) = \int_{S^1} \frac{1 - |w|^2}{|\xi - w|^2} d\mu(\xi)$$
$$(\mathcal{Q}\mu)(w) = \int_{S^1} \mathrm{Im}\, \frac{\xi + w}{\xi - w} d\mu(w) = \int_{S^1} \frac{2\,\mathrm{Im}\,\overline{\xi} w}{|\xi - w|^2} d\mu(\xi)$$

The radial limits of these harmonic functions exist for a.e. $\xi \in S^1$ with respect to Lebesgue measure λ, and the decomposition of the measure μ into absolutely continuous and singular parts with respect to Lebesgue measure $\mu = f d\lambda + \mu_s$ can be recovered from these radial limits as follows:

Theorem 2.1. *(Fatou). For Lebesgue-a.e. $\xi \in S^1$. the radial limit of $(\mathcal{P}\mu)(w)$ exists and equals $f(\xi)$.*

Theorem 2.2. *(Poltoratski* [Pol96]*). Let Q denote the radial limit of $\mathcal{Q}\mu$ on S^1. As $t \to +\infty$, the measures $\frac{\pi}{2} t 1_{\{|Q|>t\}} d\lambda$ converge weakly to μ_s.*

Since we will be dealing with functions defined in $\mathbb{D}_\infty$, we will refer to the holomorphic function in $\mathbb{D}_\infty$ defined by $H(z) = \int_{S^1} \frac{\xi + 1/z}{\xi - 1/z} d\mu(\xi)$ as the Herglotz transform of μ on $\mathbb{D}_\infty$. Then $H(z) = \tilde{H}(w)$, where $w = 1/z$ and $\tilde{H} = \mathcal{H}\mu$ is the Herglotz transform defined above which is holomorphic in $\mathbb{D}$. Appropriate versions of the above theorems hold for the boundary values of the function H. Let the boundary values of $H, \tilde{H}$ be $P + iQ, \tilde{P} + i\tilde{Q}$ respectively. Then $P = \tilde{P} \circ r, Q = \tilde{Q} \circ r$, so for $\mu = f d\lambda + \mu_s$ and $\xi \in S^1$ we have

$$f \circ r = \tilde{P} \circ r = P$$

and

$$\begin{aligned} r^*\mu_s &= \lim_{t\to+\infty} r^*\left(\frac{\pi}{2}t1_{\{|\tilde{Q}|>t\}}d\lambda\right) \\ &= \lim_{t\to+\infty}\left(\frac{\pi}{2}t1_{\{|\tilde{Q}\circ r|>t\}}d(r^*\lambda)\right) \\ &= \lim_{t\to+\infty}\left(\frac{\pi}{2}t1_{\{|Q|>t\}}d\lambda\right) \end{aligned}$$

weakly.

3. The Loewner equation

A *hull* is a connected, full, non-trivial compact K in $\mathbb{C}$ containing the origin. Its complement Ω in $\hat{\mathbb{C}}$ is then a simply connected domain containing ∞, so there is a conformal map from the complement of the closed unit disk, $\phi : \mathbb{D}_\infty \to \Omega = \hat{\mathbb{C}} - K$ such that $\phi(\infty) = \infty$, which is unique when normalized to satisfy $\phi'(\infty) > 0$. The map ϕ has an expansion around $z = \infty$ of the form

$$\phi(z) = e^{c(K)}z + a_0 + \frac{a_1}{z} + \dots$$

where the real number $c(K)$ is called the *logarithmic capacity* of the hull K (note that the closed disk of radius R then has logarithmic capacity $\log R$).

Given a strictly increasing family of hulls $(K_t)_{-\infty<t\leq t_0}$, if the domains $\Omega_t = \hat{\mathbb{C}} - K_t$ are continuous for the Caratheodory topology (which holds if the family of hulls is Hausdorff continuous for example) and $K_t \to \{0\}$ as $t \to -\infty$, then one can continuously reparametrize the family by logarithmic capacity so that the associated conformal maps $\phi_t : \mathbb{D}_\infty \to \Omega_t$ satisfy

$$\phi_t(z) = e^t z + a_0(t) + \frac{a_1(t)}{z} + \dots$$

The family of conformal maps (ϕ_t) is called a *Loewner chain*.

For each compact $K \subset \mathbb{D}_\infty$, for each z in K the map $t \mapsto \phi_t(z)$ is C_K-Lipschitz in t for some constant C_K only depending on K ([Pom75], §6.1). Thus for each z the derivative of $\phi_t(z)$ with respect to t exists for all t outside some null set E_z, but in fact one can choose a null set E independent of z such that the derivative with respect to t exists for all z in $\mathbb{D}_\infty$. Moreover this derivative is of the form

$$\frac{d^+}{ds}_{|s=t}(\phi_t^{-1}\circ\phi_s)(z) = zH_t(z)$$

where H_t is a holomorphic function on $\mathbb{D}_\infty$ satisfying $\operatorname{Re} H_t > 0, H_t(\infty) = 1$, and hence can be written as the Herglotz transform on $\mathbb{D}_\infty$ of a unique probability measure μ_t on S^1,

$$H_t(z) = \int_{S^1}\frac{\xi + 1/z}{\xi - 1/z}d\mu_t(\xi)$$

The measures (μ_t) are called the driving or Loewner measures for the Loewner chain (ϕ_t).

4. Conformal measures of analytic circle diffeomorphisms

A conformal measure of dimension δ for a holomorphic map f is a finite measure μ on a f-invariant compact K such that

$$\mu(f(E)) = \int_E |f'(z)|^\delta d\mu(z)$$

for all measurable sets $E \subset K$, or, when f is a diffeomorphism,

$$f^*\mu = |f'|^\delta d\mu$$

where $f^*\mu$ is defined by $(f^*\mu)(E) = \mu(f(E))$. Conformal measures have been studied extensively for rational maps and Kleinian groups (as Patterson-Sullivan measures), where existence is usually proved by means of transfer operators for hyperbolic dynamical systems. While the machinery of transfer operators is not available for circle diffeomorphisms, Douady and Yoccoz prove ([DY99]) nonetheless the existence and uniqueness of s-conformal measures $\mu_{s,g}$ for any C^2 circle diffeomorphism g with irrational rotation number for every $s \in \mathbb{R}$. Moreover the measures $\mu_{s,g}$ depend continuously on s, g for the weak topology on measures and the C^1 topology on circle diffeomorphisms.

5. The semigroup $(\Phi_t)_{t\geq 0}$ on $\mathrm{Diff}^\omega_\alpha(S^1)$

Given an analytic circle diffeomorphism g with irrational rotation number α, for $\epsilon > 0$ small, there is a unique, strictly increasing, Hausdorff continuous family of Herman compacts $(A_t)_{0\leq t<\epsilon}$ totally invariant under g such that the hulls $\overline{\mathbb{D}} \cup A_t$ have logarithmic capacity t. Let $(\Omega_t)_{0\leq t<\epsilon}$ be the family of decreasing simply connected domains $\Omega_t = \hat{\mathbb{C}} - (\overline{\mathbb{D}} \cup A_t)$, and let $\phi_t : \mathbb{D}_\infty \to \Omega_t$ be conformal maps normalized so that $\phi_t(\infty) = \infty, \phi_t'(\infty) = e^t$. We let $\psi_t = \phi_t^{-1} : \Omega_t \to \mathbb{D}_\infty$.

The map $g_t := \psi_t \circ g \circ \psi_t^{-1}$ is holomorphic in an annulus contained in $\mathbb{D}_\infty$ with one boundary component equal to S^1, and (as shown in [PM97]) extends analytically across S^1 to give an analytic circle diffeomorphism with rotation number equal to α. This defines a map

$$\begin{aligned} \Phi_t : \mathcal{D}_t \subset \mathrm{Diff}^\omega_\alpha(S^1) &\to \mathrm{Diff}^\omega_\alpha(S^1) \\ g &\mapsto g_t \end{aligned}$$

where $\mathcal{D}_t$ is the set of g in $\mathrm{Diff}^\omega_\alpha(S^1)$ having a Herman compact A such that $\overline{\mathbb{D}} \cup A$ has logarithmic capacity t.

For g in $\mathrm{Diff}^\omega_\alpha(S^1)$, and $t \geq 0$, let $A_t(g)$ be a Herman compact for g such that $\overline{\mathbb{D}} \cup A_t(g)$ has logarithmic capacity t (if such a Herman compact exists), let $\Omega_t(g) := \hat{\mathbb{C}} - (\overline{\mathbb{D}} \cup A_t(g))$, and let $\psi_t(g) : \Omega_t(g) \to \mathbb{D}_\infty$ be the corresponding normalized conformal map.

Proposition 5.1. *For $s, t \geq 0$, $\Phi_t(\mathcal{D}_{s+t}) \subset \mathcal{D}_s$, and $\Phi_s \circ \Phi_t = \Phi_{s+t}$ on $\mathcal{D}_{t+s}$.*

Proof: For $g \in \mathcal{D}_{s+t}$ where $s, t \geq 0$, $\psi_t(g)$ maps $\Omega_{s+t}(g)$ conformally to a domain Ω_1 such that the compact A_1 given by the complement of Ω_1 and its reflection in the unit circle is invariant under $\Phi_t(g)$, hence $\Omega_1 = \Omega_{s'}(\Phi_t(g))$ for some $s' \geq 0$. The map $\psi_{s'}(\Phi_t(g))$ maps Ω_1 conformally to $\mathbb{D}_\infty$, so the composition $\psi := \psi_{s'}(\Phi_t(g)) \circ \psi_t(g)$ maps $\Omega_{s+t}(g)$ conformally to $\mathbb{D}_\infty$ and satisfies the normalizations $\psi(\infty) = \infty, \psi'(\infty) = e^{s'} e^t = e^{s'+t} > 0$. It follows from uniqueness of the normalized conformal mapping that $\psi = \psi_{s+t}(g)$, so $e^{s'+t} = e^{s+t}$ and hence $s' = s$. Thus $\psi_s(\Phi_t(g)) \circ \psi_t(g) = \psi_{s+t}(g)$, from which it follows that $(\Phi_s \circ \Phi_t)(g) = \Phi_{s+t}(g)$. $\diamond$

6. Infinitesimal generator of the semigroup

Let g be an analytic circle diffeomorphism with irrational rotation number α. Let $(A_t)_{0 \leq t < \epsilon}$ be the family of Herman compacts of g, and $(\mathcal{H}_t = \overline{\mathbb{D}} \cup A_t)_{0 \leq t < \epsilon}$ the associated family of hulls. Let $\Omega_t = \hat{\mathbb{C}} - \mathcal{H}_t$, and let $\phi_t : \mathbb{D}_\infty \to \Omega_t, \psi_t = \phi_t^{-1}$ be normalized conformal mappings. Then the functions $\{(\phi_t - id)/t : 0 \leq t < \epsilon\}$ form a normal family ([Pom75], §6.1).

Lemma 6.1. *Let $\{t_n\}$ be a sequence decreasing monotonically to 0 such that $(\phi_{t_n} - id)/t_n$ converges uniformly on compacts of $\mathbb{D}_\infty$ to a holomorphic function χ. Then $(\psi_{t_n} - id)/t_n$ converges uniformly on compacts in $\mathbb{D}_\infty$ to $(-\chi)$.*

Proof: Fix a compact $K \subset \mathbb{D}_\infty$, then for n large $K \subset \Omega_{t_n}$ and the maps ψ_{t_n} are defined on K. We have $\phi_{t_n} = id + t_n\chi + o(t_n)$ uniformly on K. Then $id = \phi_{t_n} \circ \psi_{t_n}$ gives

$$id = (id + t_n\chi + o(t_n)) \circ \psi_{t_n} = \psi_{t_n} + t_n\chi \circ \psi_{t_n} + o(t_n),$$

so

$$\frac{\psi_{t_n} - id}{t_n} = -\chi \circ \psi_{t_n} + o(1)$$

uniformly on K, and the lemma follows since $\psi_{t_n} \to id$ uniformly on K as $n \to \infty$. $\diamond$

Lemma 6.2. *Let $\{t_n\}$ be a sequence decreasing monotonically to 0 such that $(\phi_{t_n} - id)/t_n$ converges uniformly on compacts of $\mathbb{D}_\infty$ to a holomorphic function χ. Then $(g_{t_n} - g)/t_n$ converges uniformly on compacts in a neighbourhood V of S^1 to a holomorphic function $\dot{g}$ given (in $V \cap \mathbb{D}_\infty$) by*

$$\dot{g} = g' \cdot \chi - \chi \circ g$$

Proof: By the previous Lemma, the equality $\psi_{t_n} \circ g = g_{t_n} \circ \psi_{t_n}$ gives

$$(id - t_n\chi + o(t_n)) \circ g = g_{t_n} \circ (id - t_n\chi + o(t_n))$$

uniformly on a circle $\{|z| = R\} \subset \mathbb{D}_\infty$, so

$$g - t_n\chi \circ g + o(t_n) = g_{t_n} - g'_{t_n} t_n\chi + o(t_n)$$

thus

$$\frac{g_{t_n} - g}{t_n} = g'_{t_n} \cdot \chi - \chi \circ g + o(1) \to g' \cdot \chi - \chi \circ g$$

uniformly on $\{|z| = R\}$. Since the maps g_{t_n}, g are circle maps, $(g_{t_n} - g)/t_n$ converges uniformly on the circle $\{|z| = 1/R\} \subset \mathbb{D}$, hence by the Maximum Principle we have uniform convergence on the closed annulus $\{1/R \leq |z| \leq R\}$. $\diamond$

Theorem 6.3. *Let $\{t_n\}$ be a sequence decreasing monotonically to 0 such that $(\phi_{t_n} - id)/t_n$ converges uniformly on compacts of $\mathbb{D}_\infty$ to a holomorphic function χ. Then $H = \chi/id$ satisfies $\operatorname{Re} H > 0, H(\infty) = 1$, and is the Herglotz transform $\mathcal{H}\mu$ on $\mathbb{D}_\infty$ of $\mu = r^*\mu_{2,g}$, where $r : S^1 \to S^1$ denotes complex conjugation and $\mu_{2,g}$ is the unique 2-conformal measure of g.*

Proof: Let $\chi_n(z) = (\phi_{t_n}(z) - z)/t_n$ and let $H_n(z) = \chi_n(z)/z$ for $|z| > 1$. Since for $t > 0$, Ω_t is a proper subdomain of $\mathbb{D}_\infty$, by the Schwarz Lemma $|\phi_t(z)/z| > 1$ for all $t > 0$. The Moebius map $w = (z-1)/(z+1)$ maps $\mathbb{D}_\infty$ conformally to the half-plane $\{\operatorname{Re} w > 0\}$, so

$$\operatorname{Re}\left(\frac{t_n z H_n(z)}{\phi_{t_n}(z) + z}\right) = \operatorname{Re}\left(\frac{\phi_{t_n}(z) - z}{\phi_{t_n}(z) + z}\right) > 0$$

and letting $n \to \infty$ it follows that (using $H_n(z) \to H(z), z/(\phi_{t_n}(z)+z) \to 1/2$) that $\operatorname{Re} H(z) \geq 0$ for all z. Moreover $H_n(\infty) = (e^{t_n} - 1)/t_n$, and H_n converges uniformly to H on compacts in $\mathbb{D}_\infty$, so $H(\infty) = 1$. If $\operatorname{Re} H(z_0) = 0$ for some z_0 then by the open mapping theorem we must have $H(z) \equiv H(z_0) \in i\mathbb{R}$, contradicting $H(\infty) = 1$, hence $\operatorname{Re} H(z) > 0$ for all z. By the Herglotz Theorem, H is the Herglotz transform on $\mathbb{D}_\infty$ of a probability measure μ on S^1. Since $r^2 = id$ it suffices to show that $r^*\mu$ is 2-conformal for g.

Let $\mu = f d\lambda + \mu_s$ be the decomposition of μ into absolutely continuous and singular parts with respect to Lebesgue measure.

By Fatou's Theorem, the radial limits of $\operatorname{Re} H$ exist for a.e. $\xi \in S^1$, and equal $f \circ r$. From Lemma 6.2 we have $\dot{g} + \chi \circ g = g' \cdot \chi$, and so

$$\frac{\dot{g}}{g}(z) + (H \circ g)(z) = \frac{z g'(z)}{g(z)} H(z)$$

By Lemma 6.2, the function $\dot{g}$ is holomorphic in a neighbourhood of S^1, and, since $g_{t_n}(\xi)$ converges to $g(\xi)$ along S^1 for ξ in S^1, we have $\operatorname{Re}(\dot{g}/g)(\xi) = 0$. Moreover $\xi g'(\xi)/g(\xi) = |g'(\xi)|$ since g is a circle map, so taking real parts in the equation above as z tends to $\xi \in S^1$ radially gives $(f \circ r \circ g)(\xi) = |g'(\xi)| f(r(\xi))$, thus

$$g^*(r^*(f d\lambda)) = g^*((f \circ r) d\lambda) = (f \circ r \circ g)|g'| d\lambda = |g'|^2 (f \circ r) d\lambda$$

thus the absolutely continuous part $r^*(f d\lambda) = (f \circ r) d\lambda$ of $r^*\mu$ is 2-conformal.

It remains to show that the singular part $\nu := r^*\mu_s$ of $r^*\mu$ is 2-conformal. Taking imaginary parts in $(\dot{g}/g)(z) + (H \circ g)(z) = \frac{z g'(z)}{g(z)} H(z)$ as z tends to a point $\xi \in S^1$ radially gives

$$(\dot{g}/g)(\xi) + i(Q \circ g)(\xi) = |g'(\xi)| i Q(\xi)$$

The function $\dot{g}/g$ is holomorphic in a neighbourhood of S^1, hence bounded on S^1, so $||Q \circ g| - |g'||Q|| < M$ for some $M > 0$.

For $t \gg 1$, let ν_t denote the measure $\frac{\pi}{2} t 1_{|Q|>t} d\lambda$, so ν_t converges weakly to ν as $t \to +\infty$.

Given $\xi_0 \in S^1$ and a small $\epsilon > 0$, let $U \subset S^1$ be a small interval around ξ_0 in S^1 such that $(1 - \epsilon) < |g'(\xi)/g'(\xi_0)| < (1 + \epsilon)$ for ξ in U. For $t \gg 1$ such that

$t/(t-M) < 1+\epsilon, t/(t+M) > 1-\epsilon$, we have

$$\begin{aligned}
\nu_t(g(U)) &= \frac{\pi}{2} t \lambda(g(U) \cap \{|Q| > t\}) \\
&= \frac{\pi}{2} t \lambda(g(\{\xi \in U, |Q(g(\xi))| > t\})) \\
&\leq \frac{\pi}{2} t(1+\epsilon)|g'(\xi_0)| \lambda(\{\xi \in U, |Q(g(\xi))| > t\}) \\
&\leq \frac{\pi}{2} t(1+\epsilon)|g'(\xi_0)| \lambda(\{\xi \in U, |g'(\xi)||Q(\xi)| > t - M\}) \\
&\leq t(1+\epsilon)|g'(\xi_0)| \frac{1}{T} \nu_T(U) \quad \left(\text{where } T = \frac{t-M}{(1+\epsilon)|g'(\xi_0)|} \right) \\
&= (1+\epsilon)^3 |g'(\xi_0)|^2 \nu_T(U)
\end{aligned}$$

Similarly we have

$$\nu_t(g(U)) \geq (1-\epsilon)^3 |g'(\xi_0)|^2 \nu_{T'}(U)$$

where $T' = \frac{t+M}{(1-\epsilon)|g'(\xi_0)|}$.

Douady and Yoccoz ([DY99]) show that if $\log |g'|$ is of bounded variation (which is the case if g is C^2) then any conformal measure for g has no atoms. In particular $\nu = r^* \mu_{2,g}$ has no atoms, hence $\nu_t(U) \to \nu(U)$ for any interval U. From the above it follows that given ϵ, there is a $\delta = \delta(\epsilon) > 0$ such that if U is any interval of length less than δ, and $\xi_0 \in U$, then

$$(1-\epsilon)^3 |g'(\xi_0)|^2 \nu(U) \leq \nu(g(U)) \leq (1+\epsilon)^3 |g'(\xi_0)|^2 \nu(U)$$

Given any interval V in S^1 and $\epsilon > 0$, let $U_1, \ldots, U_n$ be a partition of V into intervals of length less than $\delta(\epsilon)$ centered around points $\xi_1, \ldots, \xi_n$. We may assume that the variation of $|g'|^2$ on each U_i is less than ϵ. Then

$$\begin{aligned}
\nu(g(V)) &= \sum_i \nu(g(U_i)) \\
&\leq \sum_i (1+\epsilon)^3 |g'(\xi_i)|^2 \nu(U_i) \\
&\leq (1+\epsilon)^3 \sum_i \int_{U_i} (|g'(\xi)|^2 + \epsilon) d\nu(\xi) \\
&= (1+\epsilon)^3 \left(\int_V |g'(\xi)|^2 d\nu(\xi) + \epsilon \right)
\end{aligned}$$

Letting ϵ tend to 0 gives $\nu(g(V)) \leq \int_V |g'|^2 d\nu$, and similarly $\nu(g(V)) \geq \int_V |g'|^2 d\nu$. It follows that $\nu = r^* \mu_s$ is 2-conformal for g, hence $r^* \mu$ is 2-conformal for g and $\mu = r^* \mu_{2,g}$. $\diamond$

7. Proofs of main results

We can now prove the main results from the Introduction.

Proof of Theorem 1.1: Given $g \in \text{Diff}^{\omega}_{\alpha}(S^1)$, by uniqueness of the 2-conformal measure $\mu_{2,g}$ of g, it follows from Theorem 6.3 any normal limit χ of the functions $(\phi_t - t)/t, t > 0$ (where $\phi_t : \mathbb{D}_\infty \to \Omega_t$ is the normalized Riemann mapping as

before) is given by $\chi(z) = z \cdot H(z)$, where H is the Herglotz transform on $\mathbb{D}_\infty$ of $r^*\mu_{2,g}$. Since the normal limit is unique,

$$\frac{\phi_t(z) - z}{t} \to z \cdot H(z)$$

uniformly on compacts in $\mathbb{D}_\infty$ as $t \to 0$.

For any $t > 0$ and $s > t$, the map $\phi_t^{-1} \circ \phi_s$ is the normalized Riemann mapping from $\mathbb{D}_\infty$ to the complement of the unique hull of logarithmic capacity $s - t$ associated to the circle map g_t, so it follows from the same argument as above (applied to g_t) that

$$\frac{\phi_t^{-1} \circ \phi_s(z) - z}{s - t} \to z \cdot H_t(z)$$

uniformly on compacts in $\mathbb{D}_\infty$ as $s \to t$, where H_t is the Herglotz transform on $\mathbb{D}_\infty$ of $r^*\mu_{2,g_t}$. $\diamond$

Proof of Theorem 1.2: Given $g \in \mathrm{Diff}^\omega_\alpha(S^1)$ and $g_t = \Phi_t(g)$, since $(\phi_t - id)/t$ converges uniformly on compacts in $\mathbb{D}_\infty$ to a unique function χ as $t \to 0$, it follows from Lemma 6.2 that for any sequence $\{t_n\}$ converging to 0, the functions $(g_{t_n} - g)/t_n$ converge uniformly on compacts in a neighbourhood V of S^1 to a holomorphic function $\dot{g}$ on V satisfying (in $V \cap \mathbb{D}_\infty$)

$$\dot{g} = g' \cdot \chi - \chi \circ g$$

So all normal limits of the normal family $\{(g_t - g)/t\}$ coincide, thus

$$\frac{g_t - g}{t} \to g' \cdot \chi - \chi \circ g$$

as $t \to 0$, where $\chi(z) = z \cdot H(z)$, H being the Herglotz transform on $\mathbb{D}_\infty$ of $r^*\mu_{2,g}$. $\diamond$

Proof of Theorem 1.3: Let $g \in \mathrm{Diff}^\omega_\alpha(S^1)$ and let $\{g_t\}_{0 \le t < \epsilon} \subset \mathrm{Diff}^\omega_\alpha(S^1)$ be such that $g_0 = g$ and such that the right-hand derivatives exist uniformly on a neighbourhood V of S^1, and satisfy

$$\dot{g}_t := \frac{d^+}{ds}_{s=t} g_s(z) = X(g_t)(z) = g'(z) \cdot \chi_t(z) - \chi_t \circ g_t(z)$$

where $\chi_t(z) = zH_t(z)$, $H_t(z)$ being the Herglotz transform on $\mathbb{D}_\infty$ of the measure $r^*\mu_{2,g_t}$. The function $p(z,t) := H_t(z)$ satisfies $\mathrm{Re}\, p(z,t) > 0, p(\infty,t) = 1$. The map g_t depends continuously on t with respect to the topology of uniform convergence in a neighbourhood of S^1, and hence, being analytic maps, with respect to C^1 convergence on S^1, so by [DY99] the measures $r^*\mu_{2,g_t}$ depend continuously on t for the weak topology, hence $p(z,t) = H_t(z)$ depends continuously on t for fixed z. It follows ([Pom75] §6.1) that there exists a Loewner chain $(\phi_t)_{0 \le t < \epsilon}$, where the maps ϕ_t are normalized conformal mappings from $\mathbb{D}_\infty$ onto a decreasing family of simply connected domains Ω_t, such that $\phi_t(z) = e^t z + O(1)$ near $z = \infty$, and the following right-hand derivatives exist uniformly on compacts of $\mathbb{D}_\infty$:

$$\frac{d^+}{ds}_{|s=t} \phi_t^{-1} \circ \phi_s = \chi_t$$

Let $(A_t)_{0 \le t < \epsilon}$ be the increasing family of annular compacts containing S^1 given by the complement in $\hat{\mathbb{C}}$ of Ω_t and its reflection in S^1, so that the hulls $\mathcal{H}_t := \hat{\mathbb{C}} - \Omega_t$ are given by $\mathcal{H}_t = \overline{\mathbb{D}} \cup A_t$.

Let $\tilde{g}_t = \phi_t \circ g_t \circ \phi_t^{-1}$, then $\tilde{g}_t$ depends continuously on t with respect to the topology of uniform convergence in a neighbourhood of S^1, hence so does $\tilde{g}_t'$. Let $U = V \cap \mathbb{D}_\infty$ be a one-sided neighbourhood of S^1. For $s > t$, let $h = s - t$, then as $h \to 0$, we have, uniformly on any compact in U,

$$\begin{aligned}
\tilde{g}_s' &= \tilde{g}_t' + o(h), \\
\phi_s &= \phi_t \circ (id + h\chi_t + o(h)) = \phi_t + h\phi_t'\chi_t + o(h), \\
g_s &= g_t + h\dot{g}_t + o(h)
\end{aligned}$$

so $\tilde{g}_s \circ \phi_s = \phi_s \circ g_s$ gives:

$$\begin{aligned}
\tilde{g}_s \circ (\phi_t + h\phi_t'\chi_t + o(h)) &= (\phi_t + h\phi_t'\chi_t + o(h)) \circ (g_t + h\dot{g}_t + o(h)) \\
\Rightarrow \tilde{g}_s \circ \phi_t + h(\tilde{g}_s' \circ \phi_t)(\phi_t'\chi_t) + o(h) &= \phi_t \circ g_t + h(\phi_t' \circ g_t)(\chi_t \circ g_t + \dot{g}_t) + o(h) \\
\Rightarrow \tilde{g}_s \circ \phi_t + h(\tilde{g}_t' \circ \phi_t)(\phi_t'\chi_t) + o(h) &= \tilde{g}_t \circ \phi_t + h(\phi_t' \circ g_t)(g_t'\chi_t) + o(h) \\
\Rightarrow (\tilde{g}_s - \tilde{g}_t) \circ \phi_t + h(\tilde{g}_t \circ \phi_t)'\chi_t + o(h) &= h(\phi_t \circ g_t)'\chi_t + o(h) \\
\Rightarrow (\tilde{g}_s - \tilde{g}_t) \circ \phi_t + h(\tilde{g}_t \circ \phi_t)'\chi_t + o(h) &= h(\tilde{g}_t \circ \phi_t)'\chi_t + o(h)
\end{aligned}$$

from which it follows that for any z in U, the right-hand derivative $\frac{d^+}{dt}\tilde{g}_t(z) = 0$ for all t. Since $t \mapsto \tilde{g}_t(z)$ is continuous, we have $\tilde{g}_t(z) = \tilde{g}_0(z) = g(z)$ for all t.

The map $\tilde{g}_t$, being the conjugate of the circle map g_t by the map ϕ_t on U, maps the annulus $\phi_t(U)$ to the annulus $\phi_t(g_t(U))$. Both these annuli have $\partial\mathcal{H}_t$ as one boundary component, and $\tilde{g}_t = g$ extends analytically across S^1 to be univalent in a neighbourhood of S^1 containing $\mathcal{H}_t$, hence g leaves A_t invariant. It follows that A_t is the unique annular compact of g such that the hull $\mathcal{H}_t = \overline{\mathbb{D}} \cup A_t$ has logarithmic capacity t, hence $g_t = \Phi_t(g)$. $\diamond$

Proof of Theorem 1.5: Let f be a germ with rotation number α. Let $(K_t)_{-\infty<t<t_0}$ be the 1-parameter family of Siegel compacts of f parametrized by their logarithmic capacities, and let $\phi_t : \mathbb{D}_\infty \to \hat{\mathbb{C}} - K_t$ be normalized conformal mappings such that $\phi_t(\infty) = \infty, \phi_t(z) = e^t z + O(1)$ near $z = \infty$. Fix discs $\mathbb{D}_r \subset \mathbb{D}_{r_0}$ with $0 < r < r_0$ such that f maps $\mathbb{D}_r$ univalently into $\mathbb{D}_{r_0}$. Then for $t \ll -1$, the circle map g_t^f (given by conjugating f by ϕ_t^{-1}) is univalent on $\phi^{-1}(\mathbb{D}_r - K_t)$, which is an annulus in $\mathbb{D}_\infty$ with modulus tending to $+\infty$ as $t \to -\infty$. Therefore the family (g_t^f) forms a normal family, and any normal limit of g_t^f as $t \to -\infty$ must be a circle map univalent on $\mathbb{D}_\infty$, hence equal to R_α, so $g_t^f \to R_\alpha$ as $t \to -\infty$.

If two germs of integral curves $(g_t^{f_1}), (g_t^{f_2})$ are equal, then there is a $t_0 \in \mathbb{R}$ and neighbourhoods D_1, D_2 of the origin such that for $t < t_0$ there is a map h_t univalent on $\hat{\mathbb{C}} - K_t(f_1)$ with $h_t'(\infty) = 1$ conjugating f_1 on $D_1 - K_t(f_1)$ to f_2 on $D_2 - K_t(f_2)$ (where $K_t(f_i), i = 1, 2$ denotes the Siegel compact of f_i of logarithmic capacity t). The maps $(h_t)_{t<t_0}$ form a normal family, any normal limit of which is univalent on $\mathbb{C}^*$ and has derivative 1 at ∞, hence must be the identity. Thus $h_t \to id$ as $t \to -\infty$, and $f_1 = f_2$.

Finally, given a backward integral curve $(g_t)_{-\infty<t\leq c}$, let $\{t_n\}$ be a sequence in $(-\infty, c]$ decreasing to $-\infty$. Perez-Marco shows in [PM97] that for the circle map g_{t_0}, there exists a germ f_{t_0} with a Siegel compact K such that the fundamental

construction of [PM97] applied to the pair (f_{t_0}, K) gives the circle map g_{t_0}. Conjugating by a scaling if necessary, we may assume that K has logarithmic capacity t_0, so $K = K_{t_0}(f_{t_0})$ and $g_{t_0} = g_{t_0}^{f_{t_0}}$. By Theorem 1.3, for $t \in [t_0, c]$, $g_t = \Phi_{t-t_0}(g_{t_0})$, so pulling back a Herman compact for g_t to the plane of g_{t_0} and then to the plane of f_{t_0} gives a Siegel compact $K_t(f_{t_0})$ for f_{t_0} of logarithmic capacity t such that $g_t = g_t^{f_{t_0}}$.

Similarly for any $n \geq 0$ we obtain a germ f_{t_n} such that $g_t = g_t^{f_{t_n}}$ for all $t \in [t_n, c]$. So f_{t_n} is given by conjugating g_c by a conformal map $\phi_n : \mathbb{D}_\infty \to \hat{\mathbb{C}} - K_{t_n}(f_{t_n})$ with $\phi_n(z) = e^c z + O(z)$ near $z = \infty$. The maps $\{\phi_n\}$ are conformal mappings on $\mathbb{D}_\infty$ with fixed derivative at ∞ and hence form a normal family. Fix an annulus $U \subset \mathbb{D}_\infty$ with boundary components S^1 and a Jordan curve γ such that g_c is univalent on U. The maps $\{f_{t_n}\}$ are normalized univalent functions on the Jordan domains D_n bounded by $\phi_n(\gamma)$, and hence form a normal family as well (since the Jordan curves $\phi_n(\gamma)$ are equicontinuous). Any normal limit f then satisfies $g_t = g_t^f$ for all $t \leq c$. $\diamond$

The fact ([PM97]) that any circle map g arises from a pair (f, K), where f is a germ with a Siegel compact K also gives:

Proposition 7.1. *For any $g \in Diff_\alpha^\omega(S^1)$, there exists a backward integral curve $(g_t)_{-\infty<t\leq c}$ with $g_c = g$.*

Proof: Given a pair (f, K) which gives rise to the circle map g, let $(K_t)_{-\infty<t\leq c}$ be the unique family of Siegel compacts of f parametrized by logarithmic capacity, with $K_c = K$. Applying the fundamental construction of [PM97] to each pair (f, K_t) gives a backward integral curve $(g_t)_{-\infty<t\leq c}$ with $g_c = g$. $\diamond$

8. Linearizable maps and conformal radius of linearization domains

Let $g \in \mathrm{Diff}_\alpha^\omega(S^1)$ be a circle map which is analytically linearizable. For such a map we have uniqueness in both forward and backward time for any integral curve with initial condition g:

Theorem 8.1. *Let $(g_t^i)_{-\infty<t<c} \subset Diff_\alpha^\omega(S^1), i = 1, 2$ be two integral curves of X such that $g_{c_0}^1 = g_{c_0}^2 = g$ for some $c_0 < c$. Then $g_t^1 = g_t^2$ for all $t < c$.*

Proof: By Theorem 1.3 we have $g_t^1 = g_t^2$ for $c_0 \leq t < c$. For $t < c_0$, we have $g = \Phi_s(g_t^1) = \Phi_s(g_t^2)$ where $s = c_0 - t > 0$. Let $U \subset \mathbb{D}_\infty$ be an invariant annulus for g with boundary components equal to S^1 and a Jordan curve $\gamma \subset \mathbb{D}_\infty$. Let ϕ^1, ϕ^2 be the normalized conformal mappings defined on $\mathbb{D}_\infty$ conjugating g to g_t^1, g_t^2 respectively. For $i = 1, 2$, $\phi^i(\gamma)$ is an invariant Jordan curve for g_t^i in $\mathbb{D}_\infty$, and it follows that g_t^i is analytically linearizable. So the Herman compact for g_t^i which gives rise to g (on conjugating by ϕ^i) is a g_t^i-invariant annulus $A^i \subset \mathbb{D}_\infty$ with boundary components equal to S^1 and an invariant Jordan curve γ^i, such that ϕ^i maps $\mathbb{D}_\infty$ conformally to $\hat{\mathbb{C}} - (\mathbb{D} \cup A^i)$ and conjugates the action of g on S^1 to that of g_t^i on γ^i.

Let $L = \phi^2 \circ (\phi^1)^{-1} : \hat{\mathbb{C}} - (\mathbb{D} \cup A^1) \to \hat{\mathbb{C}} - (\mathbb{D} \cup A^2)$, then $L(\infty) = \infty, L'(\infty) = 1$, and L conjugates the action of g_t^1 on γ^1 to that of g_t^2 on γ^2. Fix a point $z_1 \in \gamma^1$ and

let $z_2 = L(z_1) \in \gamma^2$. For $i = 1, 2$ let η^i be a conformal map from a round annulus $\{r_i < |z| < 1\}$ to A^i mapping S^1 to γ^i such that $\eta^i(z_i) = 1$, then η^i conjugates the rotation R_α to g^i_t.

Suppose $r_1 \geq r_2$. Then the map $\nu := \eta^2 \circ (\eta^1)^{(-1)}$ maps A^1 into A_2, conjugates the action of g^1_t on γ^1 to that of g^2_t on γ^2, and $\nu(z_1) = L(z_1) = z_2$, hence $\nu = L$ on γ^1 (since the maps ν, L differ on γ^1 by post-composition with a homeomorphism of γ^2 commuting with g^2_t, which must be identity if it fixes a point of γ^2).

It follows that L extends to a univalent map from $\mathbb{D}_\infty$ into $\mathbb{D}_\infty$ (by setting $L = \nu$ on $A_1 \cup \gamma^1$). Since $L'(\infty) = 1$, the Schwarz Lemma implies that $L = id$, hence $g^1_t = g^2_t$.

A similar argument works if $r_2 \geq r_1$. $\diamond$

We recall that the *conformal radius* $r(D, z_0)$ of a simply connected domain D with a basepoint $z_0 \in D$ is defined by $r(D, z_0) = h'(0) > 0$, where $h : \mathbb{D} \to D$ is a conformal map from the unit disk to D satisfying the normalizations $h(0) = z_0, h'(0) > 0$. Note that a disc of radius R centered around z_0 has conformal radius R. If D is a simply connected domain in $\hat{\mathbb{C}}$ with basepoint $z_0 = \infty$ then the conformal radius is defined to be $r = e^{-t}$ where t is the logarithmic capacity of the hull $\hat{\mathbb{C}} - D$ (so the domain $\{|z| > R\}$ has conformal radius $1/R$).

Let f be a linearizable germ, and let $(K_t)_{-\infty<t\leq c}$ be the family of Siegel compacts of f parametrized by logarithmic capacity. Let D be the Siegel disk (maximal linearization domain) of f, and let $h : \mathbb{D} \to D$ be the normalized conformal mapping satisfying $h(0) = 0, h'(0) > 0$. For some $t_0 \leq c$, the interiors of the Siegel compacts K_t for $t \leq t_0$ are linearization domains $D_t \subset D$ for f bounded by analytic Jordan curves γ_t. Let $r(t) = r(D_t, 0)$ be the conformal radius of D_t, and let $R = r(D, 0)$ be the conformal radius of the Siegel disk. The normalized conformal mappings of the domains D_t are given by the maps $h_t : \mathbb{D} \to D_t, w \mapsto h((r(t)/R)w)$. Let $\Omega_t = \hat{\mathbb{C}} - K_t$, and let $\phi_t : \mathbb{D}_\infty \to \Omega_t$ be the normalized conformal map satisfying $\phi_t(\infty) = \infty, \phi'_t(\infty) > 0$.

Since $\gamma_t = \partial D_t = \partial \Omega_t$ is an analytic Jordan curve the maps h_t, ϕ_t extend analytically across S^1, and define an associated 'welding homeomorphism', which is the analytic circle map $w_t := h_t^{-1} \circ \phi_{t|S^1} : S^1 \to S^1$. The analytic circle map $k_t := w_t^{-1}$ conjugates the rotation R_α to the circle map $g_t = g^f_t$ (arising from the pair (f, K_t)).

Lemma 8.2. *For $t < t_0$,*

(i) The conformal radius $r = r(t)$ of the interior D_t of γ_t depends smoothly on the conformal radius e^{-t} of the exterior Ω_t of γ_t.

(ii) The map k_t depends smoothly on t.

Proof: (i) For $r \in (0, r(t_0))$, let $t = t(r) \in (-\infty, t_0)$ be the logarithmic capacity of the hull $K_t = h(\{|w| \leq r/R\})$. The parametrizations $h_{|\{|w|=r/R\}}$ of the Jordan curves γ_t depend smoothly on r, hence the boundary values $\phi_{t|S^1}$ of the normalized conformal mappings ϕ_t depend smoothly on r as well ([dCP03], Thm 3.4). Since $\phi'_t(\infty) = e^t$ is given in terms of these boundary values by Cauchy's integral formula, it follows that $t = t(r)$ depends smoothly on r. As $r \mapsto t(r)$ is strictly increasing

(by the Schwarz Lemma), the inverse mapping $t \in (-\infty, t_0) \mapsto r = r(t) \in (0, r(t_0))$ is smooth.

(ii) Since the boundary values $\phi_{t|S^1}$ depend smoothly on r and $t = t(r)$ is smooth by (i), these boundary values depend smoothly on t as well. By [dCP03], Thm 3.9, the welding maps w_t depend smoothly on t, hence so do their inverses k_t. $\diamond$

Theorem 8.3. *Let H_t be the Herglotz transform on $\mathbb{D}_\infty$ of the measure $r^*\mu_{2,g_t}$, and let $P_t + iQ_t$ be the boundary values on S^1 of H_t. Then for $\xi \in S^1$, we have*

$$(P_t \circ k_t)(\xi) = \frac{r'(t)}{r(t)} \cdot |k_t'(\xi)|$$

and

$$\frac{\dot{k_t}(\xi)}{k_t(\xi)} + iQ_t(k_t(\xi)) = 0$$

(where $\dot{k_t}(\xi)$ denotes the derivative with respect to t).

Proof: For $s = t + \epsilon$ with $\epsilon > 0$ small, and $w \in \mathbb{D}_\infty$ close to S^1, we have

$$(k_t^{-1} \circ \phi_t^{-1} \circ \phi_s \circ k_s)(w) = \frac{r(s)}{r(t)}(w)$$

As $\epsilon \to 0$, we have $\phi_t^{-1} \circ \phi_s = id + \epsilon\chi_t + o(\epsilon)$, where $\chi_t(z) = zH_t(z)$, and $k_s = k_t + \epsilon\dot{k_t} + o(\epsilon), r(s)/r(t) = 1 + \epsilon r'(t)/r(t) + o(\epsilon)$, hence

$$\begin{aligned} k_t^{-1} \circ (id + \epsilon\chi_t) \circ (k_t + \epsilon\dot{k_t})(w) + o(\epsilon) &= \left(1 + \epsilon\frac{r'(t)}{r(t)}\right) w + o(\epsilon) \\ \Rightarrow k_t^{-1} \circ (k_t + \epsilon(\dot{k_t} + \chi_t \circ k_t))(w) + o(\epsilon) &= \left(1 + \epsilon\frac{r'(t)}{r(t)}\right) w + o(\epsilon) \\ \Rightarrow \left(id + \epsilon((k^{-1})_t' \circ k_t)(\dot{k_t} + \chi_t \circ k_t)\right)(w) + o(\epsilon) &= \left(1 + \epsilon\frac{r'(t)}{r(t)}\right) w + o(\epsilon) \end{aligned}$$

thus

$$\frac{\dot{k_t}(w) + (\chi_t \circ k_t)(w)}{k_t'(w)} = \frac{r'(t)}{r(t)} w$$

from which we obtain

$$\frac{\dot{k_t}(w)}{k_t(w)} + (H_t \circ k_t)(w) = \frac{r'(t)}{r(t)} \cdot \frac{k_t'(w)}{k_t(w)} \cdot w$$

Since g_t is analytically linearizable, the measure $r^*\mu_{2,g_t}$ is absolutely continuous with respect to Lebesgue measure and has a smooth density. So for any $\xi \in S^1$, letting w tend to ξ radially in the equation above and taking real and imaginary parts gives the equalities asserted in the theorem (using the fact that $\dot{k_t}(\xi)/k_t(\xi)$ is purely imaginary and $\xi k_t'(\xi)/k_t(\xi) = |k_t'(\xi)|$ since the maps k_t are circle maps). $\diamond$

References

[Bis05] K. Biswas. Smooth combs inside hedgehogs. *DISCRETE AND CONTINUOUS DYNAMICAL SYSTEMS, vol. 12, 5*, pages 853–880, 2005.

[Bis08] K. Biswas. Hedgehogs of hausdorff dimension one. *Ergodic Theory and Dynamical Systems, vol. 28, 6*, pages 1713–1727, 2008.

[Bis15] K. Biswas. Positive area and inaccessible fixed points for hedgehogs. *Ergodic Theory and Dynamical Systems, http://dx.doi.org/10.1017/etds.2014.143*, pages 1–12, 2015.

[BPM13] K. Biswas and R. Perez-Marco. Uniformization of higher genus finite type log-riemann surfaces. *Preprint, http://arxiv.org/pdf/1305.2339.pdf*, 2013.

[BPM15a] K. Biswas and R. Perez-Marco. Caratheodory convergence of log-riemann surfaces and euler's formula. *Contemporary Mathematics, Volume 639, DOI: http://dx.doi.org/10.1090/conm/639*, pages 197–203, 2015.

[BPM15b] K. Biswas and R. Perez-Marco. Uniformization of simply connected finite type log-riemann surfaces. *Contemporary Mathematics, Volume 639, DOI: http://dx.doi.org/10.1090/conm/639*, pages 205–216, 2015.

[Brj71] A. D. Brjuno. Analytical form of differential equations. *Transactions Moscow Math. Soc. 25*, pages 131–288, 1971.

[Che11] A. Cheritat. Relatively compact siegel disks with non-locally connected boundaries. *Mathematische Annalen,* **349** *(3)*, pages 529–542, 2011.

[dCP03] M. Lanza de Cristoforis and L. Preciso. Differentiability properties of some nonlinear operators associated to the conformal welding in schauder spaces. *Hiroshima Math. J., No. 33*, pages 59–86, 2003.

[DY99] R. Douady and J. C. Yoccoz. Nombre de rotation des diffeomorphismes du dercles et mesures automorphes. *Regular and Chaotic Dynamics, Vol. 4, no. 4*, pages 3–24, 1999.

[PM93] R. Perez-Marco. Sur les dynamiques holomorphes non-linearisables et une conjecture de v.i. arnold. *Annales Scientifiques de l'E.N.S 26*, pages 565–644, 1993.

[PM94] R. Perez-Marco. Topology of julia sets and hedgehogs. *preprint, Université de Paris-Sud*, 1994.

[PM95] R. Perez-Marco. Uncountable number of symmetries for non-linearisable holomorphic dynamics. *Inventiones Mathematicae 119*, pages 67–127, 1995.

[PM96] R. Perez-Marco. Hedgehogs dynamics. *preprint, University of California Los Angeles*, 1996.

[PM97] R. Perez-Marco. Fixed points and circle maps. *Acta Mathematica 179:2*, pages 243–294, 1997.

[PM00] R. Perez-Marco. Siegel disks with smooth boundary. *preprint*, 2000.

[Pol96] A. G. Poltoratski. On the distribution of boundary values of cauchy integrals. *Proc. of the AMS, Vol. 124, No. 8*, pages 2455–2463, 1996.

[Pom75] C. Pommerenke. Univalent functions. *Vandenhoeck und Ruprecht*, 1975.

[Sie42] C. L. Siegel. Iteration of analytic functions. *Ann. Math., 43*, pages 807–812, 1942.

[Yoc95] J. C. Yoccoz. Petits diviseurs en dimension 1. *Asterisque 231*, 1995.

Ramakrishna Mission Vivekananda University, Belur Math, WB-711202, India
email: kingshook@rkmvu.ac.in